

GENERALISED PATTERN AVOIDANCE

ANDERS CLAEISSON

ABSTRACT. Recently, Babson and Steingrímsson have introduced generalised permutation patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation. We will consider pattern avoidance for such patterns, and give a complete solution for the number of permutations avoiding any single pattern of length three with exactly one adjacent pair of letters. For eight of these twelve patterns the answer is given by the Bell numbers. For the remaining four the answer is given by the Catalan numbers. We also give some results for the number of permutations avoiding two different patterns. These results relate the permutations in question to Motzkin paths, involutions and non-overlapping partitions. Furthermore, we define a new class of set partitions, called monotone partitions, and show that these partitions are in one-to-one correspondence with non-overlapping partitions.

1. INTRODUCTION

In the last decade a wealth of articles has been written on the subject of pattern avoidance, also known as the study of “restricted permutations” and “permutations with forbidden subsequences”. Classically, a pattern is a permutation $\sigma \in \mathcal{S}_k$, and a permutation $\pi \in \mathcal{S}_n$ avoids σ if there is no subsequence in π whose letters are in the same relative order as the letters of σ . For example, $\pi \in \mathcal{S}_n$ avoids 132 if there is no $1 \leq i < j < k \leq n$ such that $\pi(i) < \pi(k) < \pi(j)$. In [4] Knuth established that for all $\sigma \in \mathcal{S}_3$, the number of permutations in \mathcal{S}_n avoiding σ equals the n th Catalan number, $C_n = \frac{1}{1+n} \binom{2n}{n}$. One may also consider permutations that are required to avoid several patterns. In [5] Simion and Schmidt gave a complete solution for permutations avoiding any set of patterns of length three. Even patterns of length greater than three have been considered. For instance, West showed in [8] that permutations avoiding both 3142 and 2413 are enumerated by the Schröder numbers, $S_n = \sum_{i=0}^n \binom{2n-i}{i} C_{n-i}$.

In [1] Babson and Steingrímsson introduced generalised permutation patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation. The motivation for Babson and Steingrímsson in introducing these patterns was the study of Mahonian statistics, and they showed that essentially all Mahonian permutation statistics in the literature can be written as linear combinations of such patterns. An example of a generalised pattern is $(a-cb)$. An $(a-cb)$ -subword of a permutation $\pi = a_1 a_2 \cdots a_n$ is a subword $a_i a_j a_{j+1}$, ($i < j$), such that $a_i < a_{j+1} < a_j$. More generally, a pattern p is a word over the alphabet $a < b < c < d \cdots$ where two adjacent letters may or may not be separated by a dash. The absence of a dash between two adjacent letters in a p indicates that the corresponding letters in a p -subword of a permutation must be adjacent. Also, the ordering of the letters in the p -subword must match the ordering of the letters in the pattern. This definition, as well as any other definition in the introduction, will be stated rigorously in Section 2. All classical patterns are generalised patterns where each pair of adjacent letters is separated by a dash. For example, the generalised pattern equivalent to 132 is $(a-c-b)$.

We extend the notion of pattern avoidance by defining that a permutation avoids a (generalised) pattern p if it does not contain any p -subwords. We show that this is a fruitful extension, by establishing connections to other well known combinatorial structures, not previously shown to be related to pattern avoidance. The main results are given below.

P	$ \mathcal{S}_n(P) $	Description
$a-bc$	B_n	Partitions of $[n]$
$a-cb$	B_n	Partitions of $[n]$
$b-ac$	C_n	Dyck paths of length $2n$
$a-bc, ab-c$	B_n^*	Non-overlapping partitions of $[n]$
$a-bc, a-cb$	I_n	Involutions in \mathcal{S}_n
$a-bc, ac-b$	M_n	Motzkin paths of length n

Here $\mathcal{S}_n(P) = \{\pi \in \mathcal{S}_n : \pi \text{ avoids } p \text{ for all } p \in P\}$, and $[n] = \{1, 2, \dots, n\}$. When proving that $|\mathcal{S}_n(a-bc, ab-c)| = B_n^*$ (the n th Bessel number), we first prove that there is a one-to-one correspondence between $\{a-bc, ab-c\}$ -avoiding permutations and *monotone partitions*. A partition is monotone if its non-singleton blocks can be written in increasing order of their least element and increasing order of their greatest element, simultaneously. This new class of partitions is then shown to be in one-to-one correspondence with non-overlapping partitions.

2. PRELIMINARIES

By an *alphabet* X we mean a non-empty set. An element of X is called a *letter*. A *word* over X is a finite sequence of letters from X . We consider also the *empty word*, that is, the word with no letters; it is denoted by ϵ . Let $x = x_1x_2 \cdots x_n$ be a word over X . We call $|x| := n$ the *length* of x . A *subword* of x is a word $v = x_{i_1}x_{i_2} \cdots x_{i_k}$, where $1 \leq i_1 < i_2 < \cdots < i_k \leq n$. A *segment* of x is a word $v = x_i x_{i+1} \cdots x_{i+k}$. If X and Y are two linearly ordered alphabets, then two words $x = x_1x_2 \cdots x_n$ and $y = y_1y_2 \cdots y_n$ over X and Y , respectively, are said to be *order equivalent* if $x_i < x_j$ precisely when $y_i < y_j$.

Let $X = A \cup \{-\}$ where A is a linearly ordered alphabet. For each word x let \bar{x} be the word obtained from x by deleting all dashes in x . A word p over X is called a *pattern* if it contains no two consecutive dashes and \bar{p} has no repeated letters. By slight abuse of terminology we refer to the *length of a pattern* p as the length of \bar{p} . If the i th letter in p is a dash precisely when the i th letter in q is a dash, and p and q are order equivalent, then p and q are *equivalent*. In what follows all patterns will be over the alphabet $\{a, b, c, d, \dots\} \cup \{-\}$ where $a < b < c < d < \cdots$.

Let $[n] := \{1, 2, \dots, n\}$ (so $[0] = \emptyset$). A *permutation* of $[n]$ is bijection from $[n]$ to $[n]$. Let \mathcal{S}_n be the set of permutations of $[n]$. We shall usually think of a permutation π as the word $\pi(1)\pi(2) \cdots \pi(n)$ over the alphabet $[n]$. In particular, $\mathcal{S}_0 = \{\epsilon\}$, since there is only one bijection from \emptyset to \emptyset , the empty map. We say that a subword σ of π is a *p -subword* if by replacing (possibly empty) segments of π with dashes we can obtain a pattern q equivalent to p such that $\bar{q} = \sigma$. However, all patterns that we will consider will have a dash at the beginning and one at the end. For convenience, we therefore leave them out. For example, $(a-bc)$ is a pattern, and the permutation 491273865 contains three $(a-bc)$ -subwords, namely 127, 138, and 238. A permutation is said to be *p -avoiding* if it does not contain any p -subwords. Define $\mathcal{S}_n(p)$ to be the set of p -avoiding permutations in \mathcal{S}_n and, more generally, $\mathcal{S}_n(A) = \bigcap_{p \in A} \mathcal{S}_n(p)$.

We may think of a pattern p as a permutation statistic, that is, define $p\pi$ as the number of p -subwords in π , thus regarding p as a function from \mathcal{S}_n to \mathbb{N} . For example, $(a-bc)491273865 = 3$. In particular, π is p -avoiding if and only if $p\pi = 0$. We say that

two permutation statistics stat and stat' are *equidistributed* over $A \subseteq \mathcal{S}_n$, if

$$\sum_{\pi \in A} x^{\text{stat } \pi} = \sum_{\pi \in A} x^{\text{stat}' \pi}.$$

In particular, this definition applies to patterns.

Let $\pi = a_1 a_2 \cdots a_n \in \mathcal{S}_n$. An i such that $a_i > a_{i+1}$ is called a *descent* in π . We denote by $\text{des } \pi$ the number of descents in π . Observe that des can be defined as the pattern (ba) , that is, $\text{des } \pi = (ba)\pi$. A *left-to-right minimum* of π is an element a_i such that $a_i < a_j$ for every $j < i$. The number of left-to-right minima is a permutation statistic. Analogously we also define *left-to-right maximum*, *right-to-left minimum*, and *right-to-left maximum*.

In this paper we will relate permutations avoiding a given set of patterns to other better known combinatorial structures. Here follows a brief description of these structures. Two excellent references on combinatorial structures are [7] and [6].

Set partitions. A *partition* of a set S is a family, $\pi = \{A_1, A_2, \dots, A_k\}$, of pairwise disjoint non-empty subsets of S such that $S = \cup_i A_i$. We call A_i a *block* of π . The total number of partitions of $[n]$ is called a *Bell number* and is denoted B_n . For reference, the first few Bell numbers are

$$1, 1, 2, 5, 15, 52, 203, 877, 4140, 21147, 115975, 678570, 4213597.$$

Let $S(n, k)$ be the number of partitions of $[n]$ into k blocks; these numbers are called the *Stirling numbers of the second kind*.


Non-overlapping partitions. Two blocks A and B of a partition π *overlap* if

$$\min A < \min B < \max A < \max B.$$

A partition is *non-overlapping* if no pairs of blocks overlap. Thus

$$\pi = \{\{1, 2, 5, 13\}, \{3, 8\}, \{4, 6, 7\}, \{9\}, \{10, 11, 12\}\}$$

is non-overlapping. A pictorial representation of π is



Let B_n^* be the number of non-overlapping partitions of $[n]$; this number is called the n th *Bessel number* [3, p. 423]. The first few Bessel numbers are


$$1, 1, 2, 5, 14, 43, 143, 509, 1922, 7651, 31965, 139685, 636712.$$

We denote by $S^*(n, k)$ the number of non-overlapping partitions of $[n]$ into k blocks.

Involutions. An *involution* is a permutation which is its own inverse. We denote by I_n the number of involutions in \mathcal{S}_n . The sequence $\{I_n\}_0^\infty$ starts with

$$1, 1, 2, 4, 10, 26, 76, 232, 764, 2620, 9496, 35696, 140152.$$

Dyck paths. A *Dyck path* of length $2n$ is a lattice path from $(0, 0)$ to $(2n, 0)$ with steps $(1, 1)$ and $(1, -1)$ that never goes below the x -axis. Letting u and d represent the steps $(1, 1)$ and $(1, -1)$ respectively, we code such a path with a word over $\{u, d\}$. For example, the path




is coded by $uuduudd$. A *return step* in a Dyck path δ is a d such that $\delta = \alpha u \beta d \gamma$, for some Dyck paths α , β , and γ . A useful observation is that every non-empty Dyck path δ can be uniquely decomposed as $\delta = u \alpha d \beta$, where α and β are Dyck paths. This is the so-called *first return decomposition* of δ .

The n th *Catalan number* $C_n = \frac{1}{n+1} \binom{2n}{n}$ counts the number of Dyck paths of length $2n$. The sequence of Catalan numbers starts with

$$1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, 208012.$$

Motzkin paths. A *Motzkin path* of length n is a lattice path from $(0, 0)$ to $(n, 0)$ with steps $(1, 0)$, $(1, 1)$, and $(1, -1)$ that never goes below the x -axis. Letting ℓ , u , and d represent the steps $(1, 0)$, $(1, 1)$, and $(1, -1)$ respectively, we code such a path with a word over $\{\ell, u, d\}$. For example, the path



is coded by $u\ell\ell u d d \ell$. If δ is a non-empty Motzkin path, then δ can be decomposed as $\delta = \ell \gamma$ or $\delta = u \alpha d \beta$, where α , β and γ are Motzkin paths.

The n th *Motzkin number* M_n is the number of Motzkin paths of length n . The first few of the Motzkin numbers are

$$1, 1, 2, 4, 9, 21, 51, 127, 323, 835, 2188, 5798, 15511.$$

3. THREE CLASSES OF PATTERNS

Let $\pi = a_1 a_2 \cdots a_n \in \mathcal{S}_n$. Define the *reverse* of π as $\pi^r := a_n \cdots a_2 a_1$, and define the *complement* of π by $\pi^c(i) = n + 1 - \pi(i)$, where $i \in [n]$.

Proposition 1. *With respect to being equidistributed, the twelve pattern statistics of length three with one dash fall into the following three classes.*

- (i) $a-bc$, $c-ba$, $ab-c$, $cb-a$.
- (ii) $a-cb$, $c-ab$, $ba-c$, $bc-a$.
- (iii) $b-ac$, $b-ca$, $ac-b$, $ca-b$.

Proof. The bijections $\pi \mapsto \pi^r$, $\pi \mapsto \pi^c$, and $\pi \mapsto (\pi^r)^c$ give the equidistribution part of the result. Calculations show that these three distributions differ pairwise on \mathcal{S}_4 . \square

4. PERMUTATIONS AVOIDING A PATTERN OF CLASS ONE OR TWO

Proposition 2. *Partitions of $[n]$ are in one-to-one correspondence with $(a-bc)$ -avoiding permutations in \mathcal{S}_n . Hence $|\mathcal{S}_n(a-bc)| = B_n$.*

First proof. Recall that the Bell numbers satisfy $B_0 = 1$, and

$$B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_k.$$

We show that $|\mathcal{S}_n(a-bc)|$ satisfy the same recursion. Clearly, $\mathcal{S}_0(a-bc) = \{\epsilon\}$. For $n > 0$, let $M = \{2, 3, \dots, n+1\}$, and let S be a k element subset of M . For each $(a-bc)$ -avoiding permutation σ of S we construct a unique $(a-bc)$ -avoiding permutation π of $[n+1]$. Let τ be the word obtained by writing the elements of $M \setminus S$ in decreasing order. Define $\pi := \sigma 1 \tau$.

Conversely, if $\pi = \sigma 1 \tau$ is a given $(a-bc)$ -avoiding permutation of $[n+1]$, where $|\sigma| = k$, then the letters of τ are in decreasing order, and σ is an $(a-bc)$ -avoiding permutation of the k element set $\{2, 3, \dots, n+1\} \setminus \{i : i \text{ is a letter in } \tau\}$. \square

Second proof. Given a partition π of $[n]$, we introduce a standard representation of π by requiring that:

- (a) Each block is written with its least element first, and the rest of the elements of that block are written in decreasing order.
- (b) The blocks are written in decreasing order of their least element, and with dashes separating the blocks.

Define $\widehat{\pi}$ to be the permutation we obtain from π by writing it in standard form and erasing the dashes. We now argue that $\widehat{\pi} := a_1 a_2 \cdots a_n$ avoids $(a-bc)$. If $a_i < a_{i+1}$, then a_i and a_{i+1} are the first and the second element of some block. By the construction of $\widehat{\pi}$, a_i is a left-to-right minimum, hence there is no $j \in [i-1]$ such that $a_j < a_i$.

Conversely, π can be recovered uniquely from $\widehat{\pi}$ by inserting a dash in $\widehat{\pi}$ preceding each left-to-right minimum, apart from the first letter in $\widehat{\pi}$. Indeed, it is easy to see that the partition, π , in this way obtained is written in standard form. Thus $\pi \mapsto \widehat{\pi}$ gives the desired bijection. \square

Example. As an illustration of the map defined in the above proof, let

$$\pi = \{\{1, 3, 5\}, \{2, 6, 9\}, \{4, 7\}, \{8\}\}.$$

Its standard form is 8-47-296-153. Thus $\widehat{\pi} = 847296153$.

Proposition 3. *Let $L(\pi)$ be the number of left-to-right minima of π . Then*

$$\sum_{\pi \in \mathcal{S}_n(a-bc)} x^{L(\pi)} = \sum_{k \geq 0} S(n, k) x^k.$$

Proof. This result follows readily from the second proof of Proposition 2. We here give a different proof, which is based on the fact that the Stirling numbers of the second kind satisfy

$$S(n, k) = S(n-1, k-1) + kS(n-1, k).$$

Let $T(n, k)$ be the number of permutations in $\mathcal{S}_n(a-bc)$ with k left-to-right minima. We show that the $T(n, k)$ satisfy the same recursion as the $S(n, k)$.

Let π be an $(a-bc)$ -avoiding permutation of $[n-1]$. To insert n in π , preserving $(a-bc)$ -avoidance, we can put n in front of π or we can insert n immediately after each left-to-right minimum. Putting n in front of π creates a new left-to-right minimum, while inserting n immediately after a left-to-right minimum does not. \square

Proposition 4. *Partitions of $[n]$ are in one-to-one correspondence with $(a-cb)$ -avoiding permutations in \mathcal{S}_n . Hence $|\mathcal{S}_n(a-cb)| = B_n$.*

Proof. Let π be a partition of $[n]$. We introduce a standard representation of π by requiring that:

- (a) The elements of a block are written in increasing order.
- (b) The blocks are written in decreasing order of their least element, and with dashes separating the blocks.

(Note that this standard representation is different from the one given in the second proof of Proposition 2.) Define $\widehat{\pi}$ to be the permutation we obtain from π by writing it in standard form and erasing the dashes. It is easy to see that $\widehat{\pi}$ avoids $(a-cb)$. Conversely, π can be recovered uniquely from $\widehat{\pi}$ by inserting a dash in between each descent in $\widehat{\pi}$. \square

Example. As an illustration of the map defined in the above proof, let

$$\pi = \{\{1, 3, 5\}, \{2, 6, 9\}, \{4, 7\}, \{8\}\}.$$

Its standard form is 8-47-269-135. Thus $\widehat{\pi} = 847269135$.

Proposition 5.

$$\sum_{\pi \in \mathcal{S}_n(a-cb)} x^{1+\text{des } \pi} = \sum_{k \geq 0} S(n, k) x^k.$$

Proof. From the proof of Proposition 4 we see that π has $k + 1$ blocks precisely when $\hat{\pi}$ has k descents. \square

Proposition 6. *Involutions in \mathcal{S}_n are in one-to-one correspondence with permutations in \mathcal{S}_n that avoid $(a-bc)$ and $(a-cb)$. Hence*

$$|\mathcal{S}_n(a-bc, a-cb)| = I_n.$$

Proof. We give a combinatorial proof using a bijection that is essentially identical to the one given in the second proof of Proposition 2.

Let $\pi \in \mathcal{S}_n$ be an involution. Recall that π is an involution if and only if each cycle of π is of length one or two. We now introduce a standard form for writing π in cycle notation by requiring that:

- (a) Each cycle is written with its least element first.
- (b) The cycles are written in decreasing order of their least element.

Define $\hat{\pi}$ to be the permutation obtained from π by writing it in standard form and erasing the parentheses separating the cycles.

Observe that $\hat{\pi}$ avoids $(a-bc)$: Assume that $a_i < a_{i+1}$, that is $(a_i a_{i+1})$ is a cycle in π , then a_i is a left-to-right minimum in π . This is guaranteed by the construction of $\hat{\pi}$. Thus there is no $j < i$ such that $a_j < a_i$.

The permutation $\hat{\pi}$ also avoids $(a-cb)$: Assume that $a_i > a_{i+1}$, then a_{i+1} must be the smallest element of some cycle. Whence a_{i+1} is a left-to-right minimum in $\hat{\pi}$.

Conversely, if $\hat{\pi} := a_1 \dots a_n$ is an $\{a-bc, a-cb\}$ -avoiding permutation then the involution π is given by: $(a_i a_{i+1})$ is a cycle in π if and only if $a_i < a_{i+1}$. \square

Example. The involution $\pi = 826543719$ written in standard form is

$$(9)(7)(45)(36)(2)(18),$$

and hence $\hat{\pi} = 974536218$.

Proposition 7. *The number of permutations in $\mathcal{S}_n(a-bc, a-cb)$ with $n - k - 1$ descents equals the number of involutions in \mathcal{S}_n with $n - 2k$ fixed points.*

Proof. Under the bijection $\pi \mapsto \hat{\pi}$ in the proof of Proposition 6, a cycle of length two in π corresponds to an occurrence of (ab) in $\hat{\pi}$. Hence, if π has $n - 2k$ fixed points, then $\hat{\pi}$ has $n - k - 1$ descents. \square

Corollary 8.

$$\sum_{\pi \in \mathcal{S}_n(a-bc, a-cb)} x^{1+\text{des } \pi} = \sum_{k=0}^n \binom{n}{k} \binom{n-k}{k} \frac{k!}{2^k} x^{n-k}.$$

Proof. Let I_n^k denote the number of involutions in \mathcal{S}_n with k fixed points. Then Proposition 7 is equivalently stated as

$$\sum_{\pi \in \mathcal{S}_n(a-bc, a-cb)} x^{1+\text{des } \pi} = \sum_{k \geq 0} I_n^{n-2k} x^{n-k}. \quad (1)$$

The result now follows from the well-known and easily to derived formula

$$I_n^k = \binom{n}{k} \binom{n-k}{r} \frac{r!}{2^r}, \quad \text{where } r = \frac{n-k}{2},$$

for $n - k$ even, with $I_n^k = 0$ for $n - k$ odd. \square

5. PERMUTATIONS AVOIDING A PATTERN OF CLASS THREE

In [4] Knuth observed that there is a one-to-one correspondence between $(b-a-c)$ -avoiding permutations and Dyck paths. For completeness and future reference we give this result as a lemma, and prove it using a bijection which rests on the first return decomposition of Dyck paths. First we need a definition. For each word $x = x_1x_2 \cdots x_n$ without repeated letters, we define the *projection* of x onto \mathcal{S}_n , which we denote $\text{proj}(x)$, by

$$\text{proj}(x) = a_1a_2 \cdots a_n, \quad \text{where } a_i = |\{j \in [n] : x_j \leq x_i\}|.$$

Equivalently, $\text{proj}(x)$ is the permutation in \mathcal{S}_n which is order equivalent to x . For example, $\text{proj}(265) = 132$.

Lemma 1. $|\mathcal{S}_n(b-a-c)| = C_n$.

Proof. Let $\pi = a_1a_2 \cdots a_n$ be a permutation of $[n]$ such that $a_k = 1$. Then π is $(b-a-c)$ -avoiding if and only if $\pi = \sigma 1\tau$, where $\sigma := a_1 \cdots a_{k-1}$ is a $(b-a-c)$ -avoiding permutation of $\{n, n-1, \dots, n-k+1\}$, and $\tau := a_{k+1} \cdots a_n$ is a $(b-a-c)$ -avoiding permutation of $\{2, 3, \dots, k\}$.

We define recursively a mapping Φ from $\mathcal{S}_n(b-a-c)$ onto the set of Dyck paths of length $2n$. If π is the empty word, then so is the Dyck path determined by π , that is, $\Phi(\epsilon) = \epsilon$. If $\pi \neq \epsilon$, then we can use the factorisation $\pi = \sigma 1\tau$ from above, and define $\Phi(\pi) = u(\Phi \circ \text{proj})(\sigma)d(\Phi \circ \text{proj})(\tau)$. It is easy to see that Φ may be inverted, and hence is a bijection. \square

Lemma 2. *A permutation avoids $(b-ac)$ if and only if it avoids $(b-a-c)$.*

Proof. The sufficiency part of the proposition is trivial. The necessity part is not difficult either. Assume that π contains a $(b-a-c)$ -subword. Then there is a segment $Bm_1 \cdots m_r$ of π , where, for some $j < r$, $m_j < B$ and $m_r > B$. Now choose the largest i such that $m_i < B$, then $m_{i+1} > B$. \square

Proposition 14. *Dyck paths of length $2n$ are in one-to-one correspondence with $(b-a-c)$ -avoiding permutations in \mathcal{S}_n . Hence*

$$|\mathcal{S}_n(b-ac)| = \frac{1}{n+1} \binom{2n}{n}.$$

Proof. Follows immediately from Lemmas 1 and 2. \square

Proposition 15. *Let $L(\pi)$ be the number of left-to-right minima of π . Then*

$$\sum_{\pi \in \mathcal{S}_n(b-ac)} x^{L(\pi)} = \sum_{k \geq 0} \frac{k}{2n-k} \binom{2n-k}{n} x^k.$$

Proof. Let $R(\delta)$ denote the number of return steps in the Dyck path δ . It is well known (see [2]) that the distribution of R over all Dyck paths of length $2n$ is the distribution we claim that L has over $\mathcal{S}_n(b-ac)$.

Let γ be a Dyck path of length $2n$, and let $\gamma = uad\beta$ be its first return decomposition. Then $R(\gamma) = 1 + R(\beta)$. Let $\pi \in \mathcal{S}_n(b-ac)$, and let $\pi = \sigma 1\tau$ be the decomposition given in the proof of Lemma 1. Then $L(\pi) = 1 + L(\sigma)$. The result now follows by induction. \square

In addition, it is easy to deduce that left-to-right minima, left-to-right maxima, right-to-left minima, and right-to-left maxima all share the same distribution over $\mathcal{S}_n(b-ac)$.

Proposition 16. *Motzkin paths of length n are in one-to-one correspondence with permutations in \mathcal{S}_n that avoid $(a-bc)$ and $(ac-b)$. Hence*

$$|\mathcal{S}_n(a-bc, ac-b)| = M_n.$$

Proof. We mimic the proof of Lemma 1. Let $\pi \in \mathcal{S}_n(a-bc, ac-b)$. Since π avoids $(ac-b)$ it also avoids $(a-c-b)$ by Lemma 2 via $\pi \mapsto (\pi^c)^r$. Thus we may write $\pi = \sigma n \tau$, where $\pi(k) = n$, σ is an $\{a-bc, ac-b\}$ -avoiding permutation of $\{n-1, n-2, \dots, n-k+1\}$, and τ is an $\{a-bc, ac-b\}$ -avoiding permutation of $[n-k]$. If $\sigma \neq \epsilon$ then $\sigma = \sigma' r$ where $r = n-k+1$, or else an $(a-bc)$ -subword would be formed with n as the 'c' in $(a-bc)$. Define a map Φ from $\mathcal{S}_n(a-bc, ac-b)$ to the set of Motzkin paths by $\Phi(\epsilon) = \epsilon$ and

$$\Phi(\pi) = \begin{cases} \ell(\Phi \circ \text{proj})(\sigma) & \text{if } \pi = n\sigma, \\ u(\Phi \circ \text{proj})(\sigma) d\Phi(\tau) & \text{if } \pi = \sigma r n \tau \text{ and } r = n-k+1. \end{cases}$$

It is routine to find the inverse of Φ . □

Example. Let us find the Motzkin path associated with the $\{a-bc, ac-b\}$ -avoiding permutation 76453281.

$$\begin{aligned} \Phi(76453281) &= u\Phi(54231)d\Phi(1) \\ &= ul\Phi(4231)d\ell \\ &= ull\Phi(231)d\ell \\ &= ullud\Phi(1)d\ell \\ &= ulludld\ell \end{aligned}$$

ACKNOWLEDGEMENT

I am greatly indebted to my advisor Einar Steingrímsson, who put his trust in me and gave me the opportunity to study mathematics on a postgraduate level. This work has benefited from his knowledge, enthusiasm and generosity.

REFERENCES

- [1] E. Babson and E. Steingrímsson. Generalized permutation patterns and a classification of the Mahonian statistics. *Sém. Lothar. Combin.*, 44:Art. B44b, 18 pp. (electronic), 2000.
- [2] E. Deutsch. Dyck path enumeration. *Discrete Math.*, 204(1-3):167–202, 1999.
- [3] P. Flajolet and R. Schott. Nonoverlapping partitions, continued fractions, Bessel functions and a divergent series. *European J. Combin.*, 11(5):421–432, 1990.
- [4] D. E. Knuth. *The art of computer programming. Vol. 1: Fundamental algorithms*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont, 1969.
- [5] R. Simion and F. W. Schmidt. Restricted permutations. *European J. Combin.*, 6(4):383–406, 1985.
- [6] N. J. A. Sloane and S. Plouffe. *The encyclopedia of integer sequences*. Academic Press Inc., San Diego, CA, 1995. Also available online: <http://www.research.att.com/njas/sequences/>.
- [7] R. P. Stanley. *Enumerative combinatorics. Vol. I*. Wadsworth & Brooks/Cole Advanced Books & Software, Monterey, CA, 1986. With a foreword by Gian-Carlo Rota.
- [8] J. West. Generating trees and the Catalan and Schröder numbers. *Discrete Math.*, 146(1-3):247–262, 1995.

MATEMATIK, CHALMERS TEKNISKA HÖGSKOLA OCH GÖTEBORGS UNIVERSITET, S-412 96 GÖTEBORG, SWEDEN

E-mail address: `claesson@math.chalmers.se`