

# Turning cycle restrictions into mesh patterns via Foata’s fundamental transformation

Anders Claesson and Henning Ulfarsson

23 October 2023

## Abstract

An adjacent  $q$ -cycle is a natural generalization of an adjacent transposition. We show that the number of adjacent  $q$ -cycles in a permutation maps to the sum of occurrences of two mesh patterns under Foata’s fundamental transformation. As a corollary we resolve Conjecture 3.14 in the paper “From Hertzprung’s problem to pattern-rewriting systems” by the first author.

Let  $q$  be a positive integer. Following Brualdi and Deutsch [2], define an *adjacent  $q$ -cycle* in a permutation as a cycle of the form

$$(i, i + 1, \dots, i + q - 1).$$

In particular, an adjacent 1-cycle is a fixed point and an adjacent 2-cycle is more commonly known as an *adjacent transposition*. Brualdi and Deutsch showed, among other things, that if  $a_q(n, k)$  is the number of permutations of  $\{1, 2, \dots, n\}$  that—when expressed as a product of disjoint cycles—have exactly  $k$  adjacent  $q$ -cycles, then

$$a_q(n, k) = \sum_{j=k}^{\lfloor n/q \rfloor} (-1)^{k+j} \binom{j}{k} \frac{(n - (q-1)j)!}{j!}.$$

Foata’s fundamental transformation [4] bijectively maps a permutations  $\pi$  with  $k$  cycles to a permutation  $\sigma$  with  $k$  left-to-right minima by writing each cycle of  $\pi$  so that its leftmost element is the smallest, sorting the cycles in descending order with respect to their first element, and reading the resulting permutation  $\sigma$  as a word from left to right. For instance,

$$\pi = 213967548 = (5, 6, 7)(4, 9, 8)(3)(1, 2) \mapsto \sigma = 567498312.$$

Consider the effect of this transformation on a fixed point (an adjacent 1-cycle) such as 3 in the permutation  $\pi$  above. If it is not the smallest element, then it is mapped to a left-to-right minimum in  $\sigma$  that is directly followed by another left-to-right minimum. In terms of mesh patterns [1] it is an occurrence of

$$s_1 = \begin{array}{|c|} \hline \text{---} \\ \hline \bullet \\ \hline \text{---} \\ \hline \bullet \\ \hline \text{---} \\ \hline \end{array}$$

Indeed, the shading to the southwest of the two points guarantees that they are left-to-right minima, while the shading in the column between the points guarantees that they are adjacent. If, on the other hand, the fixed point is the smallest element, then it gets mapped to an occurrence of  $r_1 = \begin{array}{|c|} \hline \bullet \\ \hline \end{array}$ .

Similarly, let us consider an adjacent transposition  $(i, i+1)$  in  $\pi$ . Depending on whether or not  $(i, i+1)$  is the rightmost cycle of  $\pi$  (i.e., whether or not  $i = 1$ ) it is mapped to an occurrence of one of the patterns

$$r_2 = \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array} \quad \text{or} \quad s_2 = \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array}$$

in  $\sigma$ . For instance,  $(1, 2)$  in the example permutation  $\pi$  corresponds to an occurrence of  $r_2$  in  $\sigma$ . Proceeding with adjacent 3-cycles we have

$$r_3 = \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array} \quad \text{and} \quad s_3 = \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array}$$

and the cycle  $(5, 6, 7)$  in  $\pi$  corresponds to a unique occurrence of  $s_3$  in  $\sigma$ .

It should now be clear how to define  $r_q$  and  $s_q$  for any  $q \geq 1$ , and that the following theorem is true.

**Theorem 1.** *Assume that  $\pi \mapsto \sigma$  under Foata's fundamental transformation. For any  $q \geq 1$ , the number of adjacent  $q$ -cycles in  $\pi$  is equal to the sum of the number of occurrences of  $r_q$  and  $s_q$  in  $\sigma$ .*

The following corollary provides a generating function identity conjectured by the first author [3, Conjecture 3.14].

**Corollary 2.** *We have*

$$\sum_{n \geq 0} |S_n(p)| x^n = \sum_{m \geq 0} m! \left( \frac{x}{1+x^2} \right)^m, \quad \text{where } p = \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array}$$

and  $S_n(p)$  denotes the set of permutations avoiding the mesh pattern  $p$ .

*Proof.* Let us denote the right-hand side of the conjectured identity by  $F(x)$ , and let  $A(x)$  be the generating function for the number of permutations of  $[n]$  whose disjoint cycle decompositions have no adjacent transpositions. In other words, the coefficient of  $x^n$  in  $A(x)$  is  $a_2(n, 0)$ ; these numbers form sequence A177249 in the OEIS [6]. Brualdi and Deutsch [2] have shown that  $A(x)$  satisfies the differential equation

$$x^2(1+x^2)A'(x) - (1+x^2)(1-x-x^2)A(x) + 1-x^2 = 0, \quad A(0) = 1.$$

Term-wise differentiation yields

$$\frac{d}{dx} \left( \frac{F(x)}{1+x^2} \right) = \frac{(1-x-x^2)F(x) - 1+x^2}{x^2(1+x^2)}$$

and using this it is easy to verify that  $F(x)/(1+x^2)$  satisfies the same differential equation as  $A(x)$ . It thus suffices to show that  $\sum_{n \geq 0} |S_n(p)|x^n = (1+x^2)A(x)$ , or, equivalently,

$$|S_n(p)| = a_2(n, 0) + a_2(n-2, 0) \quad \text{for } n \geq 2. \quad (1)$$

As a special case of Theorem 1 we find that Foata's fundamental transformation provides a one-to-one correspondence between permutations without adjacent transpositions and permutations that avoid  $r_2$  and  $s_2$ . By symmetry (reverse followed by inverse) we may equivalently consider permutations avoiding the two patterns

$$r'_2 = \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \bullet & \bullet \\ \hline \end{array} \quad \text{and} \quad s'_2 = \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \bullet & \bullet \\ \hline \end{array}$$

By the Shading lemma [5], the pattern  $p$  is *coincident* with the pattern  $s'_2$ , in the sense that  $S_n(p) = S_n(s'_2)$  for all  $n \geq 0$ . By conditioning on whether a permutation avoiding  $s'_2$  also avoids  $r'_2$  or contains  $r'_2$  we find that


$$S_n(p) = S_n(r'_2, s'_2) \cup (Co_n(r'_2) \cap S_n(s'_2)),$$

where the union is disjoint and  $Co_n(r'_2)$  denotes the set of permutations that contain  $r'_2$ . Note that a permutation  $\pi$  contains  $r'_2$  and avoids  $s'_2$  precisely when it is the direct sum  $\pi = 21 \oplus \sigma$  of the permutation 21 and a permutation  $\sigma$  that avoids  $r'_2$  and  $s'_2$ . This establishes equation (1) and concludes the proof.  $\square$

As above, assume that  $\pi \mapsto \sigma$  under Foata's fundamental transformation. A direct consequence of the definition of this map is that the number of cycles of  $\pi$  equals the number of left-to-right minima of  $\sigma$ , which in turn is the number of occurrences of the mesh pattern



in  $\sigma$ . We have shown that the statistic “number of fixed points” translates to the mesh pattern statistic



and more generally that the number of adjacent  $q$ -cycles translates to  $r_q + s_q$ . A fixed point  $\pi(i) = i$  of  $\pi$  is said to be *strong* if  $j < i \Rightarrow \pi(j) < \pi(i)$  and  $j > i \Rightarrow \pi(j) > \pi(i)$ ; see [7, Ex. 1.32b]. In term of mesh patterns, strong fixed points are occurrences of in  $\pi$ , and it is easy to see that they map to occurrences of in  $\sigma$ , sometimes called *skew strong fixed points* [1]. What other properties of the cycle structure of  $\pi$  can be neatly expressed in terms of occurrences of mesh patterns in  $\sigma$ ? Are the examples presented here special cases of a more general phenomena?

## Acknowledgements

This work was started at Schloss Dagstuhl (Leibniz-Zentrum für Informatik), seminar 23121, and we thank the institute and the organizers for giving us the opportunity to participate.

## References

- [1] Petter Brändén and Anders Claesson. Mesh patterns and the expansion of permutation statistics as sums of permutation patterns. *Electron. J. Combin.*, 18(2):P5, 2011.
- [2] Richard A. Brualdi and Emeric Deutsch. Adjacent  $q$ -cycles in permutations. *Ann. Comb.*, 16(2):203–213, 2012.
- [3] Anders Claesson. From Hertzsprung’s problem to pattern-rewriting systems. *Algebraic Combinatorics*, 5(6):1257–1277, 2022.
- [4] Dominique Foata. Étude algébrique de certains problèmes d’analyse combinatoire et du calcul des probabilités. *Publ. Inst. Statist. Univ. Paris*, 14:81–241, 1965.
- [5] Í. Hilmarsson, I. Jónsdóttir, S. Sigurdardóttir, L. Vidarsdóttir, and Henning Ulfarsson. Wilf-classification of mesh patterns of short length. *Electronic Journal of Combinatorics*, 22, 2015.
- [6] OEIS Foundation Inc. The Online Encyclopedia of Integer Sequences. <https://oeis.org/>, 2023.
- [7] Richard P. Stanley. *Enumerative combinatorics*, volume 1 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2 edition, 2011.